


AUTHOR QUERY FORM

	Journal: YPLAS Article Number: 2064	Please e-mail or fax your responses and any corrections to: E-mail: corrections.esch@elsevier.sps.co.in Fax: +31 2048 52799
---	--	---

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.


Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the 'Q' link to go to the location in the proof.

Location in article	Query / Remark: click on the Q link to go Please insert your reply or correction at the corresponding line in the proof
<u>Q1</u>	Please confirm that given names and surnames have been identified correctly.

Thank you for your assistance.

Highlights

► We construct a plasmid vector for regulated gene expression in pneumococcus. ► Expression of target genes is induced by maltose. ► We test plasmid vector by cloning and expressing the *gfp* reporter gene. ► The expression system is stable under repression and induction conditions.





ELSEVIER

Contents lists available at SciVerse ScienceDirect

Plasmid

journal homepage: www.elsevier.com/locate/yplas

Construction of a plasmid vector based on the pMV158 replicon for cloning and inducible gene expression in *Streptococcus pneumoniae*

José A. Ruiz-Masó, Celeste López-Aguilar, Concha Nieto, Marta Sanz, Patricia Burón, Manuel Espinosa, Gloria del Solar*

Molecular Microbiology and Infection Biology Department, Centro de Investigaciones Biológicas, Consejo Superior de Investigaciones Científicas, Ramiro de Maeztu 9, E28040 Madrid, Spain

ARTICLE INFO

Article history:

Received 21 July 2011

Accepted 12 September 2011

Available online xxxx

Communicated by Saleem Khan

Keywords:

Plasmid pMV158

Cloning vector

Regulated expression

Pneumococcal *malR* gene

ABSTRACT

We report the construction of a plasmid vector designed for regulated gene expression in *Streptococcus pneumoniae*. The new vector, pLS1ROM, is based on the replicon of the streptococcal promiscuous rolling circle replication (RCR) plasmid pMV158. We inserted the controllable promoter P_M of the *S. pneumoniae malMP* operon, followed by a multi-cloning site sequence aimed to facilitate the insertion of target genes. The expression from P_M is negatively regulated by the transcriptional repressor MalR, which is released from the DNA operator sequence by growing the cells in maltose-containing media. To get a highly regulated expression of the target gene, MalR was provided in *cis* by inserting the *malR* gene under control of the constitutive P_{tet} promoter, which in pMV158 directs expression of the *tetL* gene. To test the functionality of the system, we cloned the reporter gene *gfp* from *Aequorea victoria*, encoding the green fluorescent protein (GFP). Pneumococcal cells harboring the recombinant plasmid rendered GFP fluorescence in a maltose-dependent mode with undetectable background levels in the absence of the inducer. The new vector, pLS1ROM, exhibits full structural and segregational stability and constitutes a valuable tool for genetic manipulation and regulated gene expression in *S. pneumoniae*.

© 2011 Published by Elsevier Inc.

1. Introduction

Streptococcus pneumoniae is a Gram-positive human pathogen that causes invasive infections such as pneumonia, meningitis or sepsis. In addition, pneumococcus is the leading bacterial cause of childhood mortality due to acute respiratory infections and non-epidemic meningitis in developing countries (WHO, 2009). Increasing antibiotic resistance of pathogenic *S. pneumoniae* is still an important problem despite the use of the conjugated pneumococcal vaccine and improvements in antimicrobial therapy (Perez-Trallero et al., 2010; Williams et al., 2002). Even though *S. pneumoniae* is a naturally transformable

bacterium that is able to accept and transfer plasmids originally isolated from other species (Barany and Tomasz, 1980; Stassi et al., 1981), the occurrence of indigenous plasmids in pneumococcus seems to be very low. Most of the pneumococcal plasmids characterized so far are related to pDP1 (Smith and Guild, 1979). An exception is plasmid pSpnP1, which was identified in a multidrug-resistant strain of *S. pneumoniae* (Romero et al., 2007). Sequence analysis showed that they are rolling circle plasmids of the pC194/pUB110 family.

The availability of plasmid-based genetic tools to study the expression of native and heterologous genes in different microorganisms has a special interest for both basic research and practical applications. To date, the number of non-integrative plasmid vectors available to manipulate *S. pneumoniae* is still very poor. In fact, most of the constructs are based on the broad-host-range RCR plasmid

* Corresponding author. Fax: +34 91 5360432.

E-mail address: gdelosolar@cib.csic.es (G. del Solar).

pMV158, naturally isolated from *Streptococcus agalactiae* (Burdett, 1980), and on its derivative pLS1 (Stassi et al., 1981). In addition, the pneumococcal cryptic plasmid pRMG1 (related to pDP1) has been used to construct a shuttle vector between pneumococcus and *Escherichia coli* (Muñoz et al., 1999). Vectors based on the promiscuous pMV158 replicon have been used to clone pneumococcal genes and to analyze its heterologous expression (Espinosa et al., 1984; Lacks et al., 1986). Recently, a plasmid vector based on pLS1 has been presented as an useful genetic tool to study promoter and terminator sequences in *S. pneumoniae* and *Enterococcus faecalis* (Ruiz-Cruz et al., 2010). Plasmids pLS1RGFP (Nieto et al., 2000), pLS1GFP (Fernández de Palencia et al., 2000), pMV158GFP (Nieto and Espinosa, 2003), and pLS70GFP (Acebo et al., 2000) constitute another set of pMV158 replicon-based tools for the regulated expression of the *gfp* reporter gene. In these constructs, expression of *gfp* was controlled by the regulatory mechanism of maltosaccharides utilization in *S. pneumoniae* (Nieto et al., 1997). This system has been validated in pneumococcus (Nieto et al., 2000) and used to detect pneumococcal cells by fluorescence microscopy in culture medium (Nieto and Espinosa, 2003) or to confirm its intracellular location in granulocytes and microglia (Letiembre et al., 2005; Ribes et al., 2010).

In the present work we have constructed pLS1ROM, a new pLS1-based vector designed for cloning and regulated gene expression in *S. pneumoniae*. The presence of a multi-cloning site that helps the insertion of the target gene immediately after the maltose-inducible promoter P_M , the tight transcriptional control of P_M arising from the presence in *cis* of the MalR repressor (Nieto et al., 2000), and the minimization of the plasmid size by removal of dispensable regions are the more remarkable features of the pLS1ROM pneumococcal vector. The suitability of pLS1ROM as a regulated expression vector has been tested by cloning the *gfp* reporter gene and analyzing the fluorescence of pneumococcal cells containing the resulting recombinant plasmid pLS1ROM-GFP upon maltose induction. In contrast to the previously constructed pLS1RGFP plasmid, pLS1ROM-GFP exhibits structural and segregational stability under induction conditions, which represents a significant improvement over the prior maltose-inducible pneumococcal system.

2. Materials and methods

2.1. Bacterial strains and plasmids

S. pneumoniae cells were grown in AGCH medium (Lacks, 1968) supplemented with 0.3% sucrose and 0.2% yeast extract, with selection for resistance to erythromycin (Ery, 1 µg/ml) when required. All cultures were grown at 37 °C. The constructs were done in *S. pneumoniae* 708 (*end-1 exo-1 trt-1 hex-4 malM594*; (Lacks and Greenberg, 1977)). *S. pneumoniae* R6 (wild type; Lacks, 1968) was used for the expression of *gfp* and detection of GFP fluorescence. Other plasmids used were pLS1R (initially named as pAPM22; (Puyet et al., 1993), pCL1 (Nieto et al., 2000) and pGreenTIR (Miller and Lindow, 1997).

2.2. Plasmid DNA preparation and DNA manipulations

Pneumococcal cells were grown and transformed with plasmid DNA as previously reported (Lacks et al., 1986; Stassi et al., 1981). Plasmid DNAs purified from *S. pneumoniae* were prepared as described (del Solar et al., 1987). Restriction endonucleases, the Klenow fragment of DNA polymerase I (PolIK), and T4 DNA ligase were purchased from New England Biolabs or Roche, and were used according to the manufacturer's instructions. Restriction- or PCR-derived DNA fragments employed for cloning experiments were purified from agarose gels by phenol extraction as described (Sambrook et al., 1989). The duplex oligonucleotide MCS having 5'-protuding ends compatible with BamHI-generated ends was obtained by annealing complementary oligonucleotides MCSUP and MCSDOWN (MCSUP: 5'-GATCAAGCTTCCGCGGCCCGGGC ATGCGGCC- GCACTAGG-3'; MCSDOWN: 5'-GATCCACTAG TGCGGCCG-CATGCCCGGCCCGGGAAG-3'); protruding ends are shown underlined in both oligonucleotides), which were previously purified from gel as described (Maniatis et al., 1982). Reduction of the plasmid vector size was performed by inverse PCR (Ochman et al., 1988). Two different sets of divergent primers were employed to eliminate the P_X promoter (DeIX1: 5'-TCGATCTGAGG CCTCTGAG-3'; DeIX2: 5'-TACCTCCCTGGTTCTAATCC-3') and the *tetL* gene (Deltet1: 5'-GTCATTAGTTGGCTGGTTAC-3'; Deltet2: 5'-GGTTAATGATACGCTTCC-3'), by using pLS1ROXM-MCS and pLS1ROMteLL-MCS as template, respectively. After amplification with the Phusion polymerase (Finnzymes), the resultant DNA fragments were gel-purified and incubated with T4 DNA ligase to generate circular molecules that were used to transform *S. pneumoniae* cells. To allow ligation-sealing of the cyclized DNA molecules, the primers were phosphorylated at their 5' end previously to being used in the amplification reaction. All new plasmid constructs were confirmed by automated sequencing.

2.3. Determination of structural and segregational stability

Plasmid structural and segregational stability was determined in exponentially growing cultures of *S. pneumoniae* R6 containing either pLS1ROM or pLS1ROM-GFP. Pneumococcal cells were grown under different conditions (0.3% sucrose or maltose as carbon source; presence or absence of selective pressure (Ery 1 µg/ml) for the resident plasmid). Samples were taken after the indicated number of generations in order to analyze the integrity of the plasmid DNA (by electrophoresis of total DNA preparations in 1% agarose gels; del Solar and Espinosa, 1992; Lacks et al., 1986), as well as the percentage of plasmid-containing (and hence Ery resistant) cells.

2.4. Measurement of GFP fluorescence

To measure the GFP fluorescence inside the pneumococcal cells we followed the procedure described in (Nieto et al., 2000). Pneumococcal cells harboring pLS1ROM-GFP were grown in sucrose-containing media, with Ery resistance (Ery^R) selection, to an OD₆₅₀ of 0.5 (about 4 × 10⁸ c.f.u./ml)

and then diluted 1:1000 into medium containing maltose, with or without selective pressure, and allowed to again reach an OD₆₅₀ of 0.5. Cells (1 ml from each culture) were sedimented by centrifugation and suspended in the same volume of PBS buffer (10 mM Na₂HPO₄, 1 mM KH₂PO₄, 140 mM NaCl, 3 mM KCl), pH 7.2. Aliquots (200 µl) were used to measure the fluorescence intensity on a Varioskan Flash spectral scanning multimode reader (Thermo Scientific) by excitation at a wavelength of 488 nm and detection of emission at 510 nm. As a control to determine the background fluorescence, cells harboring the plasmid vector pLS1ROM (lacking the *gfp* gene) were used.

3. Results and discussion

3.1. Construction of the regulated expression vector pLS1ROM

To construct the pLS1ROM vector we started with plasmid pLS1R, which has the pMV158 replicon and contains the *malR* gene cloned under the control of the P_{tet} promoter that directs constitutive expression of the tetracycline resistance *tetL* gene of pMV158 (Puyet et al., 1993) (Fig. 1). DNA from pLS1R was digested with HindIII and EcoRI, and treated with PollK before purifying the larger fragment. To obtain a DNA region containing the operators/promoters P_X and P_M we used plasmid pCL1 (Nieto et al., 2000). DNA from this plasmid was digested with Sall, treated with PollK, and then digested with SspI. The resultant ~1 kb fragment was purified and ligated to the pLS1R larger fragment. The ligation mixture was used to transform *S. pneumoniae*, and transformants containing the resultant construction pLS1ROXM (Fig. 1) were selected for Ery^R. Transcription initiated from P_M is oriented in

opposite direction to the other protein-encoding genes of pLS1ROXM (Fig. 1). Next step in the construction of the cloning vector was to introduce a multi-cloning site sequence (MCS) immediately after promoter P_M. To this end, the DNA from pLS1ROXM was linearized with BamHI and ligated to the 41-bp duplex oligonucleotide MCS, whose sequence had been designed to reconstitute a BamHI site at only one of its ends upon ligation with BamHI-generated fragments. Hence, the reconstituted BamHI site, which is to remain unique in the new construct pLS1ROXM-MCS, constitutes an additional single restriction site within the MCS. Moreover, the XbaI single site located between the P_M promoter and the MCS increases the number of restriction sites that can be used for cloning target genes (Fig. 1). Plasmid pLS1ROXM-MCS was obtained after transforming the ligation mixture into *S. pneumoniae* and selecting for Ery^R.

In order to improve the plasmid pLS1ROXM-MCS as an expression vector several modifications were carried out. First, we eliminated the P_X/O_X region, which is unnecessary for the MalR-mediated repression of P_M. In fact, the transcription rate from P_M has been reported to increase when the region encompassing P_X is deleted (Nieto et al., 2001). Region P_X/O_X was removed by inverse PCR, using the divergent phosphorylated primers DelX1 and DelX2, and DNA from pLS1ROXM-MCS as template. The amplification reaction yielded a linear DNA fragment corresponding to almost the entire vector except the P_X/O_X region. The amplified fragment was gel-purified and subjected to auto-ligation to render circular plasmids molecules that were used to transform *S. pneumoniae*. The resultant plasmid was termed pLS1ROM-tetL-MCS. Second, a ~1.4 kb DNA fragment containing the *tetL* gene of pMV158 was eliminated. This gene was expressed neither in pLS1 nor in pLS1ROM-tetL-MCS, as it was




Fig. 1. Regulated expression vector pLS1ROM. (A) Construction of pLS1ROM. The various steps in the construction of pLS1ROM are indicated. PMV158-born genes (black) and the *ermAM* gene (gray) are indicated by arrows. Chromosomal-born *malR* gene and the region containing the promoters P_M and P_X are indicated by open arrows and open boxes respectively. The promoter sequences are indicated by triangles pointing in the direction of transcription. (B) Sequence of the plasmid vector region encompassing the inducible promoter P_M, the operator O_M and the MCS.

248 uncoupled from its promoter. To delete this sequence we
 249 followed the same strategy of inverse PCR but using the
 250 divergent phosphorylated primers Deltet1 and Deltet2,
 251 and DNA from pLS1ROMtetL-MCS as template. This deletion
 252 did not include the transcriptional terminator of the *tetL*
 253 gene, which was left to attenuate possible convergent
 254 transcription from the *ermAM* gene. The final plasmid vector
 255 pLS1ROM is 6.8 kb in size, confers Ery^R due to the presence of
 256 the *ermAM* gene, and facilitates the insertion of the target
 257 gene immediately after the regulated promoter P_M (Gen-
 258 Bank accession [JN381945](#)).

259 3.2. Functionality of pLS1ROM: cloning and expression of the 260 *gfp* gene in the regulated expression vector, and stability of the 261 vector and recombinant plasmids

262 To assess the functionality of the constructed expres-
 263 sion system, the *gfp* reporter gene from *Aequorea victoria*,
 264 encoding GFP, was fused to the promoter P_M, and the fluo-
 265 rescence emission was measured. Specifically, we have
 266 used the *gfp* cassette from plasmid pGreenTIR, which was
 267 designed for prokaryotic transcriptional fusions and con-
 268 tains a mutated *gfp* gene (harboring the S65T “red shift”
 269 and F64L “protein solubility” amino acid changes), a trans-
 270 lational enhancer and a consensus ribosome binding site,
 271 so that synthesis of GFP is enhanced. This optimized *gfp*
 272 cassette has been efficiently expressed in *S. pneumoniae*
 273 (Nieto et al., 2000) and in other Gram-positive hosts like
 274 *Bacillus subtilis* (Serrano-Heras et al., 2005), *Lactococcus*
 275 *lactis* (Fernández de Palencia et al., 2000), and *E. faecalis*
 276 (Lorenzo-Díaz and Espinosa, 2009; Ruiz-Cruz et al., 2010).
 277 Recombinant plasmid pLS1ROM-GFP was constructed by
 278 inserting the *gfp* cassette from pGreenTIR under the control
 279 of P_M. To this end, DNA from pLS1ROM was linearized with
 280 BamHI and the resultant fragment was purified. The *gfp*
 281 cassette was extracted by digesting DNA from pGreenTIR

282 with BamHI. The resultant 810 bp DNA fragment was puri-
 283 fied and ligated to the pLS1ROM fragment.

284 The stability of pLS1ROM and pLS1ROM-GFP was tested
 285 in cultures of *S. pneumoniae* R6 growing exponentially in
 286 media containing sucrose (repression conditions) and in
 287 the absence of Ery. The results showed that both plasmids
 288 were stably inherited, with no detectable appearance of
 289 Ery-sensitive plasmid-free cells over at least 100 genera-
 290 tions (Fig. 2). Moreover, analysis of the total DNA content
 291 confirmed that the plasmids analyzed were stably main-
 292 tained with no modification or rearrangement observed
 293 after 100 generations (Fig. 2).

294 Expression vectors and their recombinant derivatives
 295 harboring the gene to be expressed should ideally exhibit
 296 structural and segregational stability. It has been reported
 297 that RCR vectors for use in Gram-positive bacteria fre-
 298 quently become structurally and/or segregationally unsta-
 299 ble upon insertion of even relatively small foreign DNA
 300 fragments (Grkovic et al., 2003; Leer et al., 1992). This is
 301 shown here not to be the case with pLS1ROM-GFP, which
 302 is stably maintained in pneumococcal cells, hence proving
 303 the suitability of pLS1ROM as a cloning vector for this
 304 bacterium.

305 We also analyzed the stability of the expression from the
 306 regulated promoter P_M by following the fluorescence
 307 emitted by the GFP protein in pLS1ROM-GFP-carrying
 308 pneumococcal cells that had been grown for several
 309 generations in the absence of both Ery and maltose. To
 310 induce GFP fluorescence, pneumococcal cultures grown
 311 under the above conditions for the indicated generations
 312 (Fig. 3A) were diluted 1:1000 into medium containing
 313 maltose and allowed to reach an OD₆₅₀ of 0.5. The results
 314 showed that the fluorescence emission is maintained at
 315 similar levels in the interval of generations analyzed
 316 (Fig. 3A), indicating that the integrity and functionality of
 317 the transcriptional fusion between P_M and the *gfp* gene is




Fig. 2. Structural and segregational stability of pLS1ROM and pLS1ROM-GFP. Total DNA extracted from cultures of pLS1ROM- or pLS1ROM-GFP-harboring pneumococcal cells that had been grown for the indicated number of generations in medium lacking Ery was analyzed by 1% agarose gel electrophoresis (left panel). To facilitate the identification of the main plasmid forms, pLS1ROM and pLS1ROM-GFP DNA samples were obtained by the alkaline lysis method and loaded to the left of the corresponding total DNA extracts. OC, open circular DNA; SC, supercoiled monomeric plasmid DNA. The percentage of Ery-resistant colonies is given in the right panel.




Fig. 3. GFP fluorescence in pneumococcal cells harboring pLS1ROM-GFP. (A) Pneumococcal cells carrying pLS1ROM or pLS1ROM-GFP were grown in medium lacking both Ery and maltose for the indicated number of generations. Production of GFP was subsequently induced by growing these cells for 10 generations in the presence of maltose. Fluorescence background levels were obtained from cells grown in the absence of maltose for 10 more generations. (B) The fluorescence emitted by pneumococcal cells carrying pLS1ROM-GFP under induction (maltose) and repression (sucrose) conditions was compared with the fluorescence due to the previously reported construction pLS1RGFP (Nieto et al., 2000) under the same conditions. The fluorescence emission data of pneumococcal cells carrying the *gfp*-less plasmids pLS1R and pLS1ROM were included as a control.




Fig. 4. Structural and segregational stability of pLS1RGFP, pLS1ROM and pLS1ROM-GFP under induction conditions. Pneumococcal cultures harboring the indicated plasmids were grown to an OD_{650} of 0.5 in medium containing sucrose with Ery^R selection (generation 0), and then diluted 1:2000 into medium containing maltose, with or without Ery. After growing the cultures for the indicated number of generations total DNA preparations were analyzed by 1% agarose gel electrophoresis. As a control the same pneumococcal cultures were diluted into medium containing sucrose, with or without Ery, and subjected to the same analysis. The main plasmid forms of pLS1ROM and pLS1ROM-GFP were identified in DNA samples prepared by alkaline lysis (lanes C). OC, open circular DNA; SC, supercoiled monomeric plasmid DNA; Del, deleted plasmid DNA.

318 preserved. These results are consistent with the new
319 expression system being structurally and segregationally
320 stable even in the absence of selective pressure, as demon-
321 strated by the results shown in Fig. 2.

322 Since transcription from the regulated promoter P_M
323 converges on and might collide with that of the *ermAM*
324 gene in both pLS1ROM (Fig. 1) and pLS1ROM-GFP (the
325 cloned *gfp* gene lacked an intrinsic terminator), it was
326 interesting to know whether, under induction conditions,
327 expression of *ermAM* was affected so that selectable levels
328 of Ery^R could not be achieved. To test this, we monitored
329 the growth rate of pneumococcal cells carrying pLS1ROM
330 or pLS1ROM-GFP under induction conditions (maltose)
331 with or without Ery^R selection. As a control, the same cells
332 were grown in medium with sucrose (repression condi-
333 tions), with or without Ery. The results (not shown) demon-
334 strated that the duplication time of the pneumococcal
335 cells grown in the presence of Ery was not significantly af-
336 fected by the presence or absence of maltose, indicating
337 that induction of transcription from P_M did not severely af-
338 fect expression of the *ermAM* gene.

339 When analyzing the expression of *gfp* from the regulated
340 promoter P_M , we observed that the fluorescence emitted in
341 induced cultures of pneumococcal cells carrying pLS1ROM-
342 GFP was remarkably higher than that of pneumococci
343 harboring the previously-constructed plasmid pLS1RGFP
344 (Fig. 3B). Furthermore, the cells carrying pLS1RGFP took
345 approximately four times longer to duplicate in maltose-
346 containing medium than in sucrose-containing medium,
347 irrespective of the presence or absence of the antibiotic
348 (not shown). Inhibition of the bacterial cell growth under
349 induction conditions was accompanied by progressive plas-
350 mid loss, close to 100% after 11 generations, in the absence
351 of Ery, or by deletions reducing significantly the plasmid
352 size in the presence of the antibiotic (Fig. 4). In contrast,
353 the cells harboring pLS1ROM-GFP did not show any
354 increase in their duplication time and the plasmid was
355 stably maintained, at least during the generations analyzed,
356 when grown in maltose-containing medium (Fig. 4). The
357 instability of pLS1RGFP in induced pneumococcal cultures
358 can explain the lower level of fluorescence observed in cells
359 carrying this plasmid compared with those harboring
360 pLS1ROM-GFP. The stability difference between these two
361 pMV158 replicon-based plasmids may arise from promoter
362 P_M directing transcription toward the replication origin in
363 pLS1RGFP (Nieto et al., 2000), which could destabilize the
364 plasmid under induction conditions. In contrast,
365 pLS1ROM-GFP has P_M orientated in opposite direction
366 relative to the plasmid replicon, and lacks the divergent
367 P_X promoter.

368 4. Conclusions

369 In this manuscript we describe the construction of a
370 new regulated expression vector, pLS1ROM, which is based
371 on the streptococcal pMV158 replicon and has been
372 specially designed to clone and express both native and
373 heterologous genes in *S. pneumoniae*. We have included a
374 MCS sequence to facilitate the insertion of the target gene,
375 improved the expression from the regulated promoter P_M

and reduced the plasmid size to increase the stability of
the vector. We have proved the suitability of this vector
by cloning and willfully expressing the *gfp* gene. In conclu-
sion, pLS1ROM is a stable and functional expression vector
that increases the spectrum of genetic tools available to
manipulate *S. pneumoniae*.

Acknowledgments

This study was supported by the Spanish Ministerio de
Ciencia e Innovación (Grants CSD2008-00013, INTERMODS
to ME; BFU2007-63575 and BFU2010-19597, PNEUMOTALK
to GdS) and the European Union (Grant EU-CP223111,
CAREPNEUMO to ME).

References

- Acebo, P. et al., 2000. Quantitative detection of *Streptococcus pneumoniae* cells harbouring single or multiple copies of the gene encoding the green fluorescent protein. *Microbiology* 146, 1267–1273.
- Barany, F., Tomasz, A., 1980. Genetic transformation of *Streptococcus pneumoniae* by heterologous plasmid deoxyribonucleic acid. *Journal of Bacteriology* 144, 698–709.
- Burdett, V., 1980. Identification of tetracycline-resistant R-plasmids in *Streptococcus agalactiae* (group B). *Antimicrobial Agents and Chemotherapy* 18, 753–760.
- del Solar, G. et al., 1987. Replication of the streptococcal plasmid pMV158 and derivatives in cell-free extracts of *Escherichia coli*. *Molecular and General Genetics* 206, 428–435.
- del Solar, G., Espinosa, M., 1992. The copy number of plasmid pLS1 is regulated by two trans-acting plasmid products: the antisense RNA II and the repressor protein, RepA. *Molecular Microbiology* 6, 83–94.
- Espinosa, M. et al., 1984. Transfer and expression of recombinant plasmids carrying pneumococcal mal genes in *Bacillus subtilis*. *Gene* 28, 301–310.
- Fernández de Palencia, P. et al., 2000. Expression of green fluorescent protein in *Lactococcus lactis*. *FEMS Microbiology Letters* 183, 229–234.
- Grkovic, S. et al., 2003. Stable low-copy-number *Staphylococcus aureus* shuttle vectors. *Microbiology* 149, 785–794.
- Lacks, S.A., 1968. Genetic regulation of maltosaccharide utilization in *pneumococcus*. *Genetics* 60, 685–706.
- Lacks, S.A., Greenberg, B., 1977. Complementary specificity of restriction endonucleases of *Diplococcus pneumoniae* with respect to DNA methylation. *Journal of Molecular Biology* 114, 153–168.
- Lacks, S.A. et al., 1986. Identification and analysis of genes for tetracycline resistance and replication functions in the broad-host-range plasmid pLS1. *Journal of Molecular Biology* 192, 753–765.
- Leer, R.J. et al., 1992. Structural and functional analysis of two cryptic plasmids from *Lactobacillus pentosus* MD353 and *Lactobacillus plantarum* ATCC 8014. *Molecular and General Genetics* 234, 265–274.
- Letiembre, M. et al., 2005. Toll-Like receptor 2 deficiency delays pneumococcal phagocytosis and impairs oxidative killing by granulocytes. *Infection and Immunity* 73, 8397–8401.
- Lorenzo-Díaz, F., Espinosa, M., 2009. Large-scale filter mating assay for intra- and inter-specific conjugal transfer of the promiscuous plasmid pMV158 in Gram-positive bacteria. *Plasmid* 61, 65–70.
- Maniatis, T., Fritsch, E.F., Sambrook, J., 1982. *Molecular cloning A laboratory manual*. Cold Spring Harbor Laboratory Press, New York.
- Miller, W.G., Lindow, S.E., 1997. An improved GFP cloning cassette designed for prokaryotic transcriptional fusions. *Gene* 191, 149–153.
- Muñoz, R. et al., 1999. Construction of a new *Streptococcus pneumoniae*-*Escherichia coli* shuttle vector based on the replicon of an indigenous pneumococcal cryptic plasmid. *International Microbiology* 2, 23–28.
- Nieto, C., Espinosa, M., 2003. Construction of the mobilizable plasmid pMV158GFP, a derivative of pMV158 that carries the gene encoding the green fluorescent protein. *Plasmid* 49, 281–285.
- Nieto, C. et al., 1997. The maltose/maltodextrin regulon of *Streptococcus pneumoniae*. *Journal of Biological Chemistry* 272, 30860–30865.
- Nieto, C. et al., 2000. Construction of a tightly regulated plasmid vector for *Streptococcus pneumoniae*: controlled expression of the green fluorescent protein. *Plasmid* 43, 205.
- Nieto, C. et al., 2001. MalR-mediated regulation of the *Streptococcus pneumoniae* malMP operon at promoter PM. *Journal of Biological Chemistry* 276, 14946–14954.

- Ochman, H. et al., 1988. Genetic applications of an inverse polymerase chain reaction. *Genetics* 120, 621–623. 446
- Perez-Trallero, E. et al., 2010. Antimicrobial resistance among respiratory pathogens in Spain: latest data and changes over 11 Years (1996–1997 to 2006–2007). *Antimicrobial Agents and Chemotherapy* 54, 2953–2959. 447
- Puyet, A. et al., 1993. Characterization of the *Streptococcus pneumoniae* maltosaccharide regulator MalR, a member of the LacI-GalR family of repressors displaying distinctive genetic features. *Journal of Biological Chemistry* 268, 25402–25408. 448
- Ribes, S. et al., 2010. Toll-Like receptor stimulation enhances phagocytosis and intracellular killing of nonencapsulated and encapsulated *Streptococcus pneumoniae* by murine microglia. *Infection and Immunity* 78, 865–871. 449
- Romero, P. et al., 2007. Isolation and characterization of a new plasmid pSpnP1 from a multidrug-resistant clone of *Streptococcus pneumoniae*. *Plasmid* 58, 51–60. 450
- Ruiz-Cruz, S. et al., 2010. Novel plasmid-based genetic tools for the study of promoters and terminators in *Streptococcus pneumoniae* and *Enterococcus faecalis*. *Journal of Microbiol Methods* 83, 156–163. 451
- Sambrook, J. et al., 1989. *Molecular cloning. A laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, New York. 452
- Serrano-Heras, G. et al., 2005. A new plasmid vector for regulated gene expression in *Bacillus subtilis*. *Plasmid* 54, 278–282. 453
- Smith, M.D., Guild, W.R., 1979. A plasmid in *Streptococcus pneumoniae*. *Journal of Bacteriology* 137, 735–739. 454
- Stassi, D.L. et al., 1981. Cloning of chromosomal genes in *Streptococcus pneumoniae*. *Proceedings of the National Academy of Sciences of the United States of America* 78, 7028–7032. 455
- WHO, 2009. Initiative for Vaccine Research (IVR). *Acute Respiratory Infections (Update September 2009): Streptococcus pneumoniae*. 456
- Williams, B.G. et al., 2002. Estimates of world-wide distribution of child deaths from acute respiratory infections. *The Lancet Infectious Diseases* 2, 25–32. 457

463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480